### health know-how

These statistics from the American College of Gastroenterology show the number of people who, at any given time in their lives, suffer from digestive ailments.

Heartbu rn Irritable bowel syndrome

60 million 53 million

**Ulcers** 20 million

Doctors estimate that one in three of us are plagued with digestive problems every day. Luckily, treating stomach disorders is easier than ever before

ew people talk about it, yet more than half the U.S. population have stomach problems. You'd think doctors' offices would be jammed with patients seeking relief, but they're not.

"Even severe heartburn sufferers tend to just live with their symptoms," says Seymour Katz, M.D., spokesperson for the American College of Gastroenterology.

The good news is no one has to put up with stomach pain anymore—researchers have made huge strides in diagnosing and treating tummy troubles, and they've unearthed great home remedies, as well.



### **HEARTBURN**

As many as 60 million Americans get that all-too-familiar burning sensation one or more times each month. If stomach

backing up into your esophagus, here's what to do:

Sleep on your left People who sleep on their left side have half as many flareups as right-sided sleepers. That's because sleeping on your left side makes the stomach slope in such a way as to allow food and acid to flow more quickly into the small intestine.

Steer clear of acidproducers People who have just two drinks with dinner have 10 times the stomach acid in their esophagus than teetotalers. Also worth avoiding: fatty foods, chocolate, caffeine and carbonated drinks they all boost acid production and loosen the valve between the stomach and the esophagus.

Shed a few pounds "If you're overweight, getting trim is one of the best heartburn remedies," says Ray E. Clouse, M.D., gastroenterologist at Washington University School of Medicine in St. Louis. "Once people lose the extra weight around their bellies, they usually can eat the foods that used to cause trouble."

Get help If you're munching over-thecounter antacids more than twice a week, see your doctor. "You could be suffering from gastroesophageal reflux disease (GERD)," says Dr. Katz. This severe form of heartburn is caused by an all-out malfunctioning of the valve between the stomach and esophagus. GERD must be treated or it can lead to a permanent narrowing of the esophagus, hoarseness and a chronic cough.

At some point in their lives, at least 20 million Americans develop ulcers in their stomach or duodenum —the first part of the small intestine. Though most ulcers are fairly small they c a n cause an astonish-

ing amount of gnawing pain—often between meals or in the middle of the night. Here's the best plan of attack:

First, beat the bugs The top cause of ulcers isn't stress—It's H. pylori bacteria that love to burrow into the mucous lining of the intestinal tract. A simple breath test can now diagnose H. pylori and a few weeks on antibiotics cures most sufferers

Say NO to NSAIDs Aspirin and nonsteroidal anti-inflammatory drugs (NSAIDs) are the second most common ulcer trigger. "They damage the lining of the stomach and duodenum, and slow the formation of the hormones needed for tissue repair," says pharmacist Jerry Hickey, R.Ph., of the Natural Pharmacy Society. If you use painkillers regularly, ask your doctor about switching to less irritating acetaminophen or low-dose corticosteroids.


# INTESTINAL GAS

First, the facts:
it's perfectly normal—although
mortifying—to
pass gas up to
20 times a
day. But if a
quick count

shows it happens to you more than that, here's what to do:

Follow a low-air diet "Seventy percent of intestinal gas is actually swallowed air," says Sheldon Zinberg, M.D., professor of gastroenterology at UCLA, Irvine. To avoid this, try eating more slowly, and give up common air-swallowing habits like sucking hard candies, chewing gum and sipping carbonated drinks.

Nix the milkshake
Lactose intolerance—
trouble breaking down milk
sugars—is another cause of
gas because undigested lactose is a prime source of food
for gas-producing bacteria.
If going cold turkey doesn't
appeal to you, try lactose-free
milk products—or take Lactaid, an enzyme supplement
that helps digest lactose.



bloating, nausea and gas. What is known is that IBS doesn't increase your risk of colon cancer or other intestinal trouble. To get your bowels back on track, start with the anti-gas remedies, then:

appeal to you, try lactose-free milk products—or take Lactaid, an enzyme supplement that helps digest lactose.

Soothe the stress
One of the key triggers of IBS flare-ups is stress. Experiment with relaxation train-

## The European Peppermint pills

Peppermint breaks up gas in the intestines and soothes crampy spasms in the colon—but in the stomach, it can trigger heartburn. The European solution: entericcoated peppermint cap-

sules. "They dissolve in the intestines, so there's no risk of heartburn," says pharmacist Jerry Hickey, R.Ph. Peppermint capsules are available in most health-food stores. Take as directed.

ing, meditation, exercise, extra sleep and delegating chores until you find the stress level your intestines can tolerate.

WHAT ELSE COULD IT BE?

Intestinal troubles that come on suddenly—or are persistent—

should be checked by a doctor. Here are some possible, treatable, causes:

IF YOU HAVE THESE SYMPTOMS...

abdominal pain, occasional fever or an intestinal

Cramping, abdominal pain that becomes constant

and disabling, fever, nausea, tenderness on the left

Severe abdominal pain that radiates through to the

Diarrhea, abdominal cramps, bloating, gas, fatigue

Pain in the lower right abdomen, fever, nausea and

Severe abdominal pain, sometimes jaundice -

Diarrhea, rectal bleeding, cramping or sharp

obstruction

side of the abdomen

and weight loss

back, or nausea and vomiting

yellowing of the skin and eyes


**IRRITABLE** 

**BOWEL** 

Keep a diary of symptoms "What triggers an IBS flare-up is different for each person," says Dr. Jones. Keep track of what you eat—and how it affects your intestines—for at least two weeks, and you'll soon know which foods to avoid.

Switch sweeteners Sorbitol, a sweetener used in many low-cal snacks, is a carbohydrate. "It passes straight to the large intestine, where it causes gas, bloating and diarrhea.

# **MOTILITY DISORDERS**



sult is bloating, feelings of fullness early in a meal, indigestion and, sometimes, nausea and vomiting. "No one knows why, but almost 80 percent of motility-disorder patients are women," says Robert S. Fisher, M.D., chief of gastroenterology at Philadelphia's Temple University School of Medicine. For athome relief, try these doctorrecommended remedies:

IT COULD BE...

Ulcerative colitis or Crohn's dis-

the bowel

pancreas

the intestines

the appendix

ease-inflammatory conditions of

Diverticulitis-small sac-like swel-

Pancreatitis—an inflammation of the

Gallstones-clumps of cholesterol or

pigment lodged in the gallbladder

Giardiasis—a parasitic infection of

Appendicitis—an inflammation of

lings in the wall of the colon

Mini-meal it One of the big motility-disorder hassles is that the stomach is too slow to empty. "So opt for five or six small, light meals each day instead of three large ones," says Dr. Fisher.

Sweat to reduce bloating More than 80 percent of women can reduce bloating simply by working out for at least two hours each week. "The human body is designed to be active," says pharmacist Jerry Hickey. "So daily exercise is one of the best ways to get your digestive tract functioning properly."

Try teas that are tummy-friendly Ginger and slippery elm teas help food move through the digestive tract smoothly. If you don't like herbal teas, try ginger or slippery elm supplements—you'll find them at your local health-food store.

od store. —Brenda Kearns ♀

12/14/98 **First** for women 43